

Pacific County Diabetes Statistics, 2003



Diabetes Prevention & Control Program
P.O. Box 47836
Olympia, WA 98504-7836
(360) 236-3617

Prevalence (adults 18+ years)	Pacific County		Washington State	
	Percent (95% CI) ¹	Number of adults	Percent (95% CI) ¹	Number of adults
	7.8 (5.2, 11.5)	1,813	6.6 (6.2, 7.1)	303,283

Risk Factors (adults 18+ years without diabetes)	Pacific County	Washington State
	Percent (95%CI) ¹	Percent (95%CI) ¹
Overweight or obese ²	70.3 (63.0, 76.7)	56.8 (55.8, 57.8)
Insufficient physical activity ³	39.1 (31.6, 47.2)	35.8 (34.9, 36.8)
Smoke cigarettes	28.0 (21.8, 35.3)	19.8 (19.0, 20.6)
History of high blood pressure ⁴	35.5 (29.0, 42.7)	20.8 (20.1, 21.5)
History of high cholesterol ⁴	41.1 (33.5, 49.2)	31.1 (30.2, 32.1)

Hospitalizations⁵ (with complications of diabetes)	Number of hospitalizations	Age-adjusted rate per 10,000 total population (95% CI) ¹		Crude rate per 10,000 people with diabetes	
	Pacific County	Pacific County	Washington State	Pacific County	Washington State
Diabetes as first listed diagnosis	25	9.0 (5.5, 14.5)	9.6 (9.4, 9.9)	137.0	191.3
Diabetes as any listed diagnosis	382	112.7 (101.3, 125.6)	118.4 (117.5, 119.3)	2,092.9	2,293.6
→ with coronary heart disease	127	36.3 (30.2, 44.0)	36.3 (35.8, 36.8)	695.8	693.1
→ with stroke	36	9.9 (6.9, 14.6)	10.1 (9.9, 10.4)	197.2	190.9
→ with diabetic ketoacidosis	7	2.6 (0.9, 6.5)	3.9 (3.8, 4.1)	38.4	78.9
→ with lower extremity amputation	8	2.2 (0.9, 5.6)	1.9 (1.8, 2.0)	43.8	160.7

Deaths⁵	Number of deaths	Age-adjusted rate per 100,000 total population (95% CI) ¹		Crude rate per 100,000 people with diabetes	
	Pacific County	Pacific County	Washington State	Pacific County	Washington State
Diabetes as underlying cause	11	34.4 (16.3, 73.2)	26.0 (24.7, 27.4)	602.7	494.4
Diabetes as any cause	40	115.9 (82.1, 167.3)	77.1 (74.9, 79.4)	2,191.5	1,460.9

¹Data presented as: percentage or rate (95% confidence interval).

²Overweight=body mass index (BMI) 25.0-29.9 kg/m², obese=BMI 30.0+ kg/m².

³Insufficient physical activity=reported level of physical activity during leisure time or at work does not meet Centers for Disease Control and Prevention (CDC) recommendations of moderate-intensity activity for 30+ minutes on 5+ days/week or vigorous-intensity activity for 20+ minutes on 3+ days/week

⁴Told by doctor, nurse, or other health professional had high blood pressure and/or high cholesterol.

⁵Rates are not computed for counts <5. Rates based on fewer than 20 deaths or hospitalizations are likely to be unstable or imprecise.